

Opioid Risk Tool - Revised (ORT-R)

The revised ORT has clinical usefulness in providing clinicians a simple, validated method to rapidly screen for the risk of developing OUD in patients on or being considered for opioid therapy.

Opioid Risk Tool – OUD (ORT-OUD)

This tool should be administered to patients upon an initial visit prior to beginning or continuing opioid therapy for pain management. A score of 2 or lower indicates low risk for future opioid use disorder; a score of ≥ 3 indicates high risk for opioid use disorder.

Mark Each Box That Applies	Yes	No
Family history of substance abuse		
Alcohol	1	0
Illegal drugs	1	0
Rx drugs	1	0
Personal history of substance abuse		
Alcohol	1	0
Illegal drugs	1	0
Rx drugs	1	0
Age between 16-45 years	1	0
Psychological disease		
ADD, OCD, bipolar, schizophrenia	1	0
Depression	1	0
Scoring total		

Cheatle, M, Compton, P, Dhingra, L, Wasser, T, O'Brien, C. (2019) Development of the Revised Opioid Risk Tool to Predict Opioid Use Disorder in Patients with Chronic Nonmalignant Pain The Journal of Pain 0 (0) 1-10. Available online: [https://www.jpain.org/article/S1526-5900\(18\)30622-9/fulltext](https://www.jpain.org/article/S1526-5900(18)30622-9/fulltext) Accessed June 10, 2019.